YR5 FRACTIONS KNOWLEDGE ORGANISER

Key Concepts


- compare and order fractions whose denominators are all multiples of the same number
- identify, name and write equivalent fractions of a given fraction, represented visually, including tenths and hundredths
- recognise mixed numbers and improper fractions and convert from one form to the other and write mathematical statements > 1 as a mixed number
- add and subtract fractions with the same denominator and denominators that are multiples of the same number
- multiply proper fractions and mixed numbers by whole numbers

Key Vocabulary

- numerator
- denominator
- equivalent
- mixed number
- improper fraction

Equivalent Fractions

Equivalent fractions have different numerators and denominators but share the same value.



If you multiply or divide the numerator and denominator of a fraction by the same number, the new fraction will be equivalent.



Improper Fractions and Mixed Numbers

An **improper fraction** has a numerator which is greater than the denominator. For example:

3 13 28 9 2 5 10

A **mixed number** is made up of an integer and a proper fraction. For example:






To **convert between improper fractions and mixed numbers**, we need to look at how many parts make up the whole.

The bar models show $\frac{13}{6}$. There are 6 parts in the whole. $13 \div 6 = 2$ remainder 1 $\frac{13}{6} = 2\frac{1}{6}$	
The bar models show $3\frac{2}{5}$.	
There are 5 parts in the whole. $3 \times 5 = 15$	
$\frac{15}{5} + \frac{2}{5} = \frac{17}{5}$	

Compare and Order Fractions

To **compare** and **order** fractions, we need to find a common denominator or numerator.






© Copyright Deepening Understanding LTD 2019

Photocopiable for educational purposes only


YR5 FRACTIONS KNOWLEDGE ORGANISER


These fractions have been ordered from smallest to greatest. Their equivalent fractions using common denominators are shown beneath.



Add Fractions

When we **add fractions** with **different denominators**, we need to find a common denominator.






Remember, when we have found the common denominator, we only need to add the numerators. We can use this method to **add three fractions** b e y o n d 1 .


3	12	10	3	. 4	5	12	, 5
7	+ 21 +	14	7	+ 7	+ 7 =	7	7

To **add mixed numbers**, we add the wholes then



Subtract Fractions

To subtract fractions with different denominators, we again find a common denominator. We can convert mixed numbers to improper fractions when we need to exchange.





© Copyright Deepening Understanding LTD 2019

Photocopiable for educational purposes only

Converting mixed numbers to proper fractions also helps us when we **subtract mixed numbers** where exchanging is needed.


	1	1	7		2		1	7		22		17		5
2-	5	- 1	10	-	2 10	-	1	10	_	10	-	10	=	10

Multiply Fractions by Integers

To **multiply a fraction by an integer**, we multiply the numerator by the integer.

3	0	6	
7 ×	2	= 7	

To **multiply a mixed number by an integer**, we can multiply the whole and part separately or convert to an improper fraction.



Fractions as Operators

We can **multiply fractions by integers** to find **fractions of amounts**.

